

## Fatty acid utilization during the early larval stages of Florida pompano (*Trachinotus carolinus*) and Common snook (*Centropomus undecimalis*)

Marion R Hauville<sup>1,2</sup>, Kevan L Main<sup>2</sup>, Hervé Migaud<sup>1</sup> & J. Gordon Bell<sup>1</sup>

<sup>1</sup>Institute of Aquaculture, University of Stirling, Stirling, Scotland, FK9 4LA, UK

<sup>2</sup>Mote Marine Laboratory, Marine and Freshwater Aquaculture Research Program, Sarasota, FL, 34240, USA

**Correspondence:** M R Hauville, Institute of Aquaculture, University of Stirling, Stirling, FK9 4LA, Scotland, UK.

E-mail: mrhauville@gmail.com

### Abstract

The pattern of conservation and loss of fatty acids from the yolk sac during the endogenous feeding period and subsequent starvation was studied in pompano and snook larvae. Fundamental information on the early fatty acid dynamic and mobilization of pompano and snook larvae was collected. In both species, fatty acids were utilized as an energy source after hatching. Mono-unsaturated fatty acids were catabolized, while saturated and poly-unsaturated fatty acids were conserved. High levels of arachidonic acid (ARA) in pompano and snook eggs, as well as selective retention in the unfed larvae suggest a high dietary requirement for this fatty acid during the early stages of larval development. The effect of an ARA supplementation was therefore investigated in snook larvae at the rotifer feeding stage. The fatty acid profile of the larvae was successfully influenced to match that of wild eggs; however, no significant improvement in growth or survival was observed. Future research should be carried out over a longer period of time and include factors related to stress resistance.

**Keywords:** fatty acid, starvation, fish, larvae, arachidonic acid, enrichment

### Introduction

In the wild, fish larvae feed mainly on copepods (Hunter 1981), which provide adequate nutrition and a large variety of prey sizes essential for successful development (Van der Meer, Olsen,

Hamre & Fyhn 2008). Nevertheless, more research is needed before mass copepod production can be economically viable and hatcheries have to rely on the traditional live preys: rotifers and *Artemia* (Stottrup 2000). Despite the convenience of their production, rotifers and *Artemia* have a main drawback which resides in their poor nutritional profile (Conceição, Yúfera, Makridis, Morais & Dinis 2010). Both prey types are deficient in essential polyunsaturated fatty acids (PUFAs), which have long been identified as critical nutritional factors in marine fish larval development and survival (Watanabe, Kitajima & Fujita 1983; Sargent, McEvoy & Bell 1997). Indeed, due to their inability to elongate and desaturate linolenic acid (18:3n-3) and linoleic acid (18:2n-6) to the PUFAs eicosapentaenoic acid (20:5n-3, EPA), docosahexaenoic acid (22:6n-3, DHA) and arachidonic acid (20:4n-6, ARA), marine fish depend solely on their diet to fulfil their requirements (Sargent, McEvoy, Estévez, Bell, Bell, Henderson & Tocher 1999; Tocher 2003).

Polyunsaturated fatty acid deficiencies strongly impact growth and normal development of fish larvae (Mourete, Rodriguez, Tocher & Sargent 1993; Watanabe 1993). DHA is highly concentrated the neural and visual cell membrane and a deficiency has been shown to impact visual development (Neuringer, Anderson & Connor 1988; Bell, Batty, Dick, Fretwell, Navarro & Sargent 1995; Benítez-Santana, Masuda, Juárez Carrillo, Ganuza, Valencia, Hernández-Cruz & Izquierdo 2007), skeletal development (Cahu, Zambonino Infante & Takeuchi 2003; Roo, Hernández-Cruz, Socorro, Fernández-Palacios, Montero & Izquierdo

2009) and stress and immune responses (Montero, Kalinowski, Obach, Robaina, Tort, Caballero & Izquierdo 2003; Ganga, Tort, Acerete, Montero & Izquierdo 2006). EPA and ARA are both precursors of eicosanoids, and compete for the cyclo-oxygenases and lipoxygenases that generate the local hormones prostaglandins, thromboxanes and leukotrienes that are involved in the modulation of neural transmission, hypothalamic functions and many immune functions (Bell, Ashton, Seccombe, Weitzel, Dick & Sargent 1996; Tocher, Bell & Sargent 1996; Sargent, McEvoy *et al.* 1999). Even though EPA is the main C20 PUFA in fish tissues, ARA seems to be the preferred substrate for eicosanoid production and the metabolites produced are of higher biological activity (Tocher *et al.* 1996). EPA competitively inhibits the production of eicosanoids from ARA therefore eicosanoid actions are determined by the ARA/EPA ratio in the tissues (Sargent, Bell, McEvoy, Tocher & Estévez 1999). Similarly, DHA and EPA compete for the formation of phospholipid structures with a higher biological value for DHA than EPA (Rodríguez, Pérez, Badía, Izquierdo, Fernández-Palacios & Lorenzo Hernandez 1998; Sargent, McEvoy *et al.* 1999; Glencross 2009). Determining the optimal DHA:EPA:ARA ratio has proven difficult, since changing the percentage of one PUFA alters the ratio of all three (Sargent, Bell *et al.* 1999). Considerable research has been carried out on cold and temperate water species, where a DHA/EPA ratio equal or greater than 2 has been considered adequate (Sargent, Bell *et al.* 1999). An ARA/EPA ratio close to 0.25 was found necessary for good growth and survival of gilthead sea bream *Sparus aurata* and European sea bass larvae *Dicentrarchus labrax* (Atalah, Hernández-Cruz, Ganuza, Benítez-Santana, Ganga, Roo, Montero & Izquierdo 2011; Atalah, Maria Hernandez-Cruz, Benitez-Santana, Ganga, Roo & Izquierdo 2011). It has been shown that an excess of ARA can cause malpigmentation in flatfish and an ARA/EPA ratio of 0.25 for turbot *Scophthalmus maximus*, Atlantic halibut *Hippoglossus hippoglossus* and Japanese flounder *Paralichthys olivaceus* and 0.5 for Senegalese sole *Solea senegalensis* gave the best percentage of normal pigmentation (McEvoy, Estevez, Bell, Shields, Gara & Sargent 1998; Estevez, Kaneko, Seikai, Tagawa & Tanaka 2001; Bell 2003; Villalta, Estévez, Bransden & Bell 2008). An ARA supplementation has proven beneficial in improving growth and survival in turbot (Bell, Castell *et al.*

1995; Bell, Batty *et al.* 1995; Castell, Bell, Tocher & Sargent 1994), gilthead sea bream (Koven, Barr, Lutzky, Ben-Atia, Weiss, Harel, Behrens & Tandler 2001) and striped bass *Morone saxatilis* (Harel, Gavasso, Leshin, Gubernatis & Place 2001). Bessonart, Izquierdo, Salhi, Hernández-Cruz, González and Fernández-Palacios (1999) have shown that, in gilthead sea bream, the effect of an ARA supplementation is enhanced if associated with a high DHA/EPA ratio, which highlights the necessity to consider FA proportions in addition to absolute amount (Sargent, McEvoy *et al.* 1999; Izquierdo, Socorro, Arantzamendi & Hernández-Cruz 2000).

On the other hand, little is known about the PUFA requirement of marine finfish from subtropical/tropical waters, as most of them are new emerging species in aquaculture. As a general trend, these species tend to have intermediate to high DHA and ARA levels and low EPA levels, leading to high DHA/EPA and ARA/EPA ratios (Gibson 1983; Fogerty, Evans, Ford & Kennett 1986; Faulk & Holt 2003; Ogata, Emata, Garibay & Furuita 2004; Yanes-Roca, Rhody, Nystrom & Main 2009). This higher ARA content suggests that larvae from tropical species may require more ARA than cold/temperate species to ensure normal development (Ogata *et al.* 2004; Faulk & Holt 2005).

Common snook and Florida pompano are species commonly found in the tropical and subtropical western Atlantic Ocean including the Gulf of Mexico and previous work has shown that eggs of both species have a high ARA/EPA ratio (Yanes-Roca *et al.* 2009; Main, Resley, Rhody, Nystrom, Stevens & Adams 2010). Since fish eggs contain all the essential nutrients required for the successful development of the embryo and the yolk-sac larvae, it is believed that their composition reflects the optimal first feeding diet (Heming & Buddington 1988). The study of the pattern of conservation and loss of fatty acids (FAs) from the yolk sac during the endogenous feeding period has proven a useful approach to investigate the requirements of fish larvae with the most critical FAs spared to the detriment of others catabolized as an energy source (Izquierdo 1996; Hamre, Yúfera, Ronnestad, Boglione, Conceição & Izquierdo 2013).

The aim of this study was to examine the changes occurring in the FA profiles of common snook and pompano larvae deprived of food after hatching to gain a better understanding of their FA requirements. In addition, an experiment was

carried out to investigate the effectiveness of an ARA supplementation in snook larvae at the rotifer feeding stage, to produce larvae with PUFA ratios close to that of natural eggs and to study the impact on early larval development.

## Materials and methods

### Starvation trial

Pompano broodstock were captured off of Florida's west coast and held inland in a zero-discharge recirculating system at the Mote Marine Laboratory Center for Marine and Freshwater Aquaculture Research, in Sarasota, Florida. Spontaneous spawning was obtained after photo-thermal conditioning in a 25 m<sup>3</sup> tank, and hormonal induction of mature females with sGnRH $\alpha$  (50  $\mu$ g kg<sup>-1</sup>). Fish were fed daily a 40% thread herring, 30% shrimp, 30% squid diet at 6.5% body weight during the conditioning period. Snook broodstock were collected in estuarine waters around Sarasota, Florida with seines in mid-July during the ambient spawning window. Capture and strip spawning procedures were similar to those described in Yanes-Roca *et al.* 2009.

Collected eggs from the captive pompano (fertilization rate = 54.3%) and wild snook (fertilization rate = 68.2%) broodstocks were placed into small conical tanks with aeration. Once the aeration was stopped, the buoyant fertilized eggs accumulated at the surface and the poor, unfertilized eggs were drained from the hatchery. The aeration was then turned back on and three 5 ml samples were taken and counted to estimate the egg concentration. Approximately 750 eggs were volumetrically stocked in sieves made of a 100 mm diameter PVC pipe sealed at one end with a 330  $\mu$ m mesh. The sieves were set on a grid in a 340 L water table (salinity 35 g L<sup>-1</sup>  $\pm$  1, dissolved oxygen 5 mg L<sup>-1</sup>  $\pm$  1, pH 8.5  $\pm$  0.5, temperature 27.5°C  $\pm$  1) equipped with a 25 W UV light and a 50 L min<sup>-1</sup> pump. In addition, three egg aliquots were sampled from the conical tank, drained on a 100  $\mu$ m sieve and rinsed with deionized water before storage at -70°C.

Larvae hatched after 24 h for pompano (hatch rate = 73.2%) and 18 h for snook (hatch rate = 96%). Three sieves were removed each day and the larvae anaesthetized with MS222 and carefully separated from egg casings and any dead eggs or larvae. The larvae from each sieve were

pooled and rinsed with deionized water then stored at -70°C. The trials were stopped when no living larvae were observed in the sieves, at 5 days post hatch (DPH) for pompano, and 6 DPH for snook.

### Live food trial

Two trials were run to study the effect of an ARA supplementation on snook larvae. Eggs for the first trial were obtained from captive broodstock captured off the west coast of Florida and held in a 45 m<sup>3</sup> tank at the Mote Marine Laboratory Center for Marine and Freshwater Aquaculture Research. Spawning was induced by photo-thermal conditioning and sGnRH $\alpha$  implantation (female only, 50  $\mu$ g kg<sup>-1</sup>). Fish were fed a 50% shrimp, 50% thread herring diet at 2.5% body weight every other day during the conditioning period. Fish spawned spontaneously (fertilization rate = 65.8%). Eggs for the second trial were collected in the wild in mid-August, using the technique described above (fertilization rate = 49.2%). In both trials, the eggs were stocked at 150 eggs L<sup>-1</sup> in 130 L tanks in a recirculating system equipped with drum filtration, biological filtration, ozonation, carbon filtration, UV lights and an inline heater unit (temperature 27.5  $\pm$  1°C, salinity 35  $\pm$  1 g L<sup>-1</sup>, dissolved oxygen of 6  $\pm$  1 mg L<sup>-1</sup> and pH of 8  $\pm$  0.5).

From 3 days after hatching (hatching rate: trial 1 = 69.8%, trial 2 = 47.6%), larvae were fed twice a day with enriched rotifers (5 mL<sup>-1</sup>) after shading with RotiGrow *plus* (Reed Mariculture Inc, Campbell, CA, USA) at 500 000 cells per litre.

During the first trial, three enrichments were tested in quadruplicate: Algamac 3050 (A1) (Aquafauna Bio-Marine Inc, Hawthorne, CA, USA), and two experimental formulations (Fa and Fb) (Reed Mariculture Inc) while during the second trial only two enrichments were tested in quadruplicate due to limited egg availability: Algamac 3050 (A2) and one experimental formulation (Fc) (Reed Mariculture Inc). The Algamac 3050 (flake form) and experimental formulations (oil-based) were emulsified prior to rotifers enrichment. Rotifers (*Brachionus plicatilis*) were batch cultured in six 200 L tanks at 27°C (salinity 35 g L<sup>-1</sup>) and fed concentrated *Nannochloropsis* (Nanno 3600, Reed Mariculture Inc). One tank was harvested daily and the rotifers were rinsed and transferred to 18 L buckets (one per enrichment) and enriched for 7 h, then rinsed and fed to the larvae or stored at 8°C overnight for the following morn-

ing feeding. Samples of enriched rotifers were preserved for FA analysis by sieving on a 55 µm mesh and rinsing with deionized water before storage at  $-70^{\circ}\text{C}$ .

At 1, 7, 12 and 15 DPH (end of the rotifer feeding period) for the first trial and 1, 5, 9 and 14 DPH for the second trial, 10 larvae per tank were photographed and standard length and eye diameter were measured using ImageJ (U.S. National Institutes of Health, Bethesda, Maryland, USA). At the end of each trial, all larvae were counted to determine survival and 50 larvae per tank were preserved for FA analysis. In addition to standard length and eye diameter, swim bladder inflation was also assessed at that point.

### Fatty acid analyses

Lipids were extracted according to Folch, Lees and Sloane Stanley (1957) and the FA composition was determined by gas-liquid chromatography after preparation of fatty acid methyl esters (FAMES) according to Morrison and Smith (1964). FAMES were analysed on a gas chromatograph (Shimadzu GC-2014, Shimadzu Scientific Instruments, Columbia, MD, USA) equipped with a flame ionization detector using a Phenomenex ZB-WAX plus capillary column (30 m long, 0.53 mm internal diameter, 1.0 µm thickness; Phenomenex, Torrance, CA, USA). The flow rate of the carrier gas helium was  $4\text{ mL min}^{-1}$ . Temperature was held at  $160^{\circ}\text{C}$  for 5 min then increased up to  $220^{\circ}\text{C}$  at  $3^{\circ}\text{C}$  per minute and maintained at this temperature for 30 min. Injector and detector temperatures were 250 and  $260^{\circ}\text{C}$  respectively. FAMES peaks were identified by comparison with known standards (Supelco, Inc., Bellefonte, PA, USA). During lipid extraction, tricosanoic acid (23:0) was added as an internal standard for subsequent quantification of FAs.

### Statistical analyses

Statistical analysis was performed with MINITAB<sup>®</sup> version 16.0 (Minitab Ltd., Coventry, UK). Growth and eye diameter data were compared using a General Linear Model (GLM) analysing all time and treatment interactions. Significant differences between treatment means were tested by Tukey's test ( $P < 0.05$ ). Non-homogeneous data (survival, swim bladder inflation and FA composition) were arcsine square root transformed before a one-way

ANOVA followed by a Tukey *post hoc* test ( $P < 0.05$ ). All data are presented as mean  $\pm$  SE of the mean.

## Results

### Fatty acid utilization

FA data are presented in Table 1 for pompano and Table 2 for snook. In addition the change in concentration of DHA, EPA, ARA, total saturated, total mono-saturated and total poly-unsaturated from the egg to the end of the starvation period are represented in Fig. 1 for pompano and Fig. 2 for snook.

In both pompano and snook, the four predominant FAs in the eggs and newly hatched larvae (0 DPH) were DHA, 16:0, 18:1n-9 and 16:1n-7, contributing to over 50% of total FA. For pompano DHA was the main FA throughout the experiment (4 DPH), while for snook 16:0 was the main FA in the egg and larvae up to 2 DPH when its proportion decreases, while DHA increases. At the end of the trial, DHA, 16:0 and 18:1n-9 were the predominant FAs, followed by 18:0 then ARA for pompano and ARA followed by 18:0 for snook.

In pompano, from hatching to the end of the trial period, proportions of DHA and ARA increased from  $33.7 \pm 0.5\%$  up to  $35.2 \pm 0.9\%$ , and  $3.2 \pm 0.1\%$  up to  $5.5 \pm 0.2\%$  respectively, while EPA decreased from  $4.7 \pm 0.2\%$  down to  $3.0 \pm 0.0\%$ . In snook, these FAs followed a similar trend with the proportion of DHA and ARA increasing from  $16.2 \pm 0.2\%$  up to  $25.8 \pm 0.5\%$ , and from  $6.5 \pm 0.2\%$  up to  $9.4 \pm 0.2\%$  respectively, while EPA decreased from  $3.0 \pm 0.2\%$  down to  $2.3 \pm 0.0\%$ . Accordingly, DHA/EPA and ARA/EPA ratios increased from  $7.1 \pm 0.1$  up to  $11.9 \pm 0.4$ , and  $0.7 \pm 0.0$  up to  $1.9 \pm 0.1$  respectively for pompano; and from  $5.4 \pm 0.2$  up to  $11.1 \pm 0.1$ , and  $2.2 \pm 0.1$  up to  $4.0 \pm 0.1$  respectively for snook.

Overall, the proportion of saturated FAs increased in both species between hatching and the end of the experiment, from  $22.5 \pm 0.8\%$  up to  $27.7 \pm 0.7\%$  in pompano larvae and from  $28.4 \pm 0.4\%$  up to  $32.4 \pm 0.4\%$  in snook larvae. In addition, the total proportion of n-6 increased from  $6.2 \pm 0.0\%$  up to  $8.5 \pm 0.1\%$  in pompano and from  $11.8 \pm 0.3\%$  up to  $15.6 \pm 0.9\%$  in snook. In contrast, the proportion of mono-unsaturated FAs decreased from  $22.8 \pm 0.7\%$  down to  $15.1 \pm 0.4\%$  in pompano and from  $28.9 \pm 0.2\%$  down to  $17.1 \pm 0.8\%$  in

**Table 1** Fatty acid profile of fertilized eggs and unfed pompano larvae up to 4 days post hatch (DPH). Only fatty acids contributing to at least 1% at one point are reported, all fatty acids are included in totals and ratios. Values are  $\pm$ SEM,  $n = 3$ . Letters indicate significant differences within a same row (Tukey test,  $p < 0.05$ )

% total FA	Fertilized egg	0 DPH	1 DPH	2 DPH	3 DPH	4 DPH
14:0	1.8 $\pm$ 0.0 <sup>c</sup>	2.0 $\pm$ 0.1 <sup>c</sup>	1.6 $\pm$ 0.1 <sup>bc</sup>	1.1 $\pm$ 0.1 <sup>ab</sup>	1.1 $\pm$ 0.0 <sup>a</sup>	0.8 $\pm$ 0.1 <sup>a</sup>
16:0	21.5 $\pm$ 0.1 <sup>b</sup>	17.1 $\pm$ 0.8 <sup>a</sup>	18.2 $\pm$ 0.1 <sup>a</sup>	17.7 $\pm$ 0.7 <sup>a</sup>	18.3 $\pm$ 0.1 <sup>a</sup>	17.5 $\pm$ 0.6 <sup>a</sup>
18:0	2.9 $\pm$ 0.0 <sup>a</sup>	2.5 $\pm$ 0.1 <sup>a</sup>	4.2 $\pm$ 0.4 <sup>b</sup>	5.8 $\pm$ 0.1 <sup>c</sup>	7.5 $\pm$ 0.2 <sup>d</sup>	8.3 $\pm$ 0.4 <sup>d</sup>
Total saturated	27.1 $\pm$ 0.1 <sup>bc</sup>	22.5 $\pm$ 0.8 <sup>a</sup>	24.9 $\pm$ 0.3 <sup>ab</sup>	25.7 $\pm$ 0.8 <sup>bc</sup>	28.1 $\pm$ 0.4 <sup>c</sup>	27.7 $\pm$ 0.7 <sup>bc</sup>
16:1n-7	4.4 $\pm$ 0.0 <sup>d</sup>	5.0 $\pm$ 0.1 <sup>d</sup>	4.8 $\pm$ 0.0 <sup>d</sup>	3.8 $\pm$ 0.2 <sup>c</sup>	3.0 $\pm$ 0.1 <sup>b</sup>	2.1 $\pm$ 0.1 <sup>a</sup>
18:1n-9	17.9 $\pm$ 0.2 <sup>d</sup>	15.0 $\pm$ 0.5 <sup>c</sup>	14.4 $\pm$ 0.5 <sup>c</sup>	13.7 $\pm$ 0.6 <sup>bc</sup>	12.0 $\pm$ 0.1 <sup>ab</sup>	10.3 $\pm$ 0.4 <sup>a</sup>
18:1n-7	2.7 $\pm$ 0.0 <sup>c</sup>	2.4 $\pm$ 0.0 <sup>bc</sup>	2.4 $\pm$ 0.1 <sup>bc</sup>	2.3 $\pm$ 0.1 <sup>b</sup>	2.1 $\pm$ 0.1 <sup>ab</sup>	1.8 $\pm$ 0.1 <sup>a</sup>
Total monounsaturated	25.6 $\pm$ 0.1 <sup>d</sup>	22.8 $\pm$ 0.7 <sup>c</sup>	22.0 $\pm$ 0.5 <sup>c</sup>	20.3 $\pm$ 0.9 <sup>c</sup>	17.8 $\pm$ 0.2 <sup>b</sup>	15.1 $\pm$ 0.4 <sup>a</sup>
18:2n-6	1.2 $\pm$ 0.0 <sup>b</sup>	1.3 $\pm$ 0.0 <sup>b</sup>	1.3 $\pm$ 0.1 <sup>b</sup>	1.0 $\pm$ 0.0 <sup>a</sup>	1.3 $\pm$ 0.4 <sup>b</sup>	0.9 $\pm$ 0.1 <sup>a</sup>
20:4n-6	2.4 $\pm$ 0.1 <sup>a</sup>	3.2 $\pm$ 0.1 <sup>b</sup>	3.7 $\pm$ 0.1 <sup>c</sup>	3.9 $\pm$ 0.1 <sup>c</sup>	4.7 $\pm$ 0.1 <sup>d</sup>	5.5 $\pm$ 0.2 <sup>e</sup>
20:5n-3	3.2 $\pm$ 0.0 <sup>a</sup>	4.7 $\pm$ 0.2 <sup>c</sup>	4.6 $\pm$ 0.2 <sup>c</sup>	3.9 $\pm$ 0.2 <sup>b</sup>	3.3 $\pm$ 0.1 <sup>a</sup>	3.0 $\pm$ 0.0 <sup>a</sup>
22:5n-6	0.9 $\pm$ 0.0 <sup>a</sup>	1.2 $\pm$ 0.1 <sup>ab</sup>	1.2 $\pm$ 0.1 <sup>ab</sup>	1.4 $\pm$ 0.1 <sup>b</sup>	1.4 $\pm$ 0.1 <sup>b</sup>	1.3 $\pm$ 0.0 <sup>b</sup>
22:5n-3	3.4 $\pm$ 0.0 <sup>c</sup>	3.4 $\pm$ 0.2 <sup>c</sup>	2.7 $\pm$ 0.0 <sup>b</sup>	2.6 $\pm$ 0.1 <sup>b</sup>	2.0 $\pm$ 0.0 <sup>a</sup>	1.8 $\pm$ 0.0 <sup>a</sup>
22:6n-3	28.8 $\pm$ 0.2 <sup>a</sup>	33.7 $\pm$ 0.5 <sup>b</sup>	32.0 $\pm$ 0.2 <sup>ab</sup>	33.1 $\pm$ 0.3 <sup>b</sup>	33.4 $\pm$ 0.6 <sup>b</sup>	35.2 $\pm$ 0.9 <sup>b</sup>
Total polyunsaturated	43.4 $\pm$ 0.3 <sup>a</sup>	51.1 $\pm$ 0.8 <sup>b</sup>	48.8 $\pm$ 0.5 <sup>b</sup>	49.1 $\pm$ 0.5 <sup>b</sup>	48.8 $\pm$ 0.7 <sup>b</sup>	50.4 $\pm$ 0.8 <sup>b</sup>
Total n-3	36.8 $\pm$ 0.2 <sup>a</sup>	43.2 $\pm$ 0.9 <sup>b</sup>	40.4 $\pm$ 0.4 <sup>ab</sup>	40.5 $\pm$ 0.5 <sup>ab</sup>	39.3 $\pm$ 0.8 <sup>ab</sup>	40.4 $\pm$ 0.8 <sup>ab</sup>
Total n-6	5.0 $\pm$ 0.1 <sup>a</sup>	6.2 $\pm$ 0.0 <sup>b</sup>	6.7 $\pm$ 0.2 <sup>c</sup>	7.0 $\pm$ 0.1 <sup>c</sup>	8.0 $\pm$ 0.1 <sup>d</sup>	8.5 $\pm$ 0.1 <sup>d</sup>
n-3/n-6 ratio	7.4 $\pm$ 0.1 <sup>c</sup>	7.0 $\pm$ 0.3 <sup>c</sup>	6.0 $\pm$ 0.1 <sup>b</sup>	5.8 $\pm$ 0.2 <sup>b</sup>	5.0 $\pm$ 0.2 <sup>a</sup>	4.8 $\pm$ 0.1 <sup>a</sup>
ARA/EPA	0.8 $\pm$ 0.0 <sup>a</sup>	0.7 $\pm$ 0.0 <sup>a</sup>	0.8 $\pm$ 0.0 <sup>a</sup>	1.0 $\pm$ 0.0 <sup>b</sup>	1.4 $\pm$ 0.1 <sup>c</sup>	1.9 $\pm$ 0.1 <sup>d</sup>
DHA/EPA	8.9 $\pm$ 0.1 <sup>b</sup>	7.1 $\pm$ 0.1 <sup>a</sup>	7.0 $\pm$ 0.3 <sup>a</sup>	8.5 $\pm$ 0.2 <sup>b</sup>	10.1 $\pm$ 0.2 <sup>c</sup>	11.9 $\pm$ 0.4 <sup>d</sup>
Total FA (mg FA g <sup>-1</sup> dry weight)	190.7 $\pm$ 3.0 <sup>c</sup>	181.5 $\pm$ 1.1 <sup>bc</sup>	179.4 $\pm$ 2.3 <sup>b</sup>	173.5 $\pm$ 1.2 <sup>b</sup>	147.5 $\pm$ 2.7 <sup>a</sup>	146.3 $\pm$ 0.7 <sup>a</sup>

snook larvae. The proportion of poly-unsaturated FAs and total proportion of n-3 increased from  $37.4 \pm 0.4\%$  up to  $48.7 \pm 0.7\%$ , and  $23.3 \pm 0.3\%$  up to  $31.4 \pm 0.9\%$  respectively in snook larvae, while contrastingly they did not vary significantly in pompano larvae with an average of  $49.7 \pm 0.2\%$  and  $40.8 \pm 0.7\%$  respectively. Consequently, the n-3/n-6 ratio decreased from  $7.0 \pm 0.3$  down to  $4.8 \pm 0.1$  in pompano, while it did not vary significantly in snook larvae with an average of  $1.9 \pm 0.8$ .

The total amount of FA decreased by 23% in pompano larvae with  $190.7 \pm 3$  mg in the eggs and  $146.3 \pm 0.7$  in the larvae at the end of the trial, and by 43% in snook larvae with  $193.3 \pm 3.5$  mg in the eggs and  $110.1 \pm 2.2$  mg at the end of the trial.

### Live feed trials

At the end of the trials, survival rates from hatching for the A1 and A2 larvae and the Fc larvae were similar (ranging from  $5.2 \pm 1.2\%$  to  $6.9 \pm 0.9\%$ ), significantly higher than that of Fa larvae with  $2.4 \pm 0.5\%$  and Fb larvae with  $2.1 \pm 0.4\%$  (Table 3). No significant difference was observed in the proportion of functional swim bladder in A1 and A2 larvae ( $80.2 \pm 2.1\%$  and  $77.1 \pm 3.1\%$

respectively); however, it was significantly higher compared to the other treatments that ranged from  $50.9 \pm 2.8\%$  to  $58.4 \pm 5.2\%$  (Table 3).


At the end of the first trial, standard length was significantly higher for the Fa and Fb larvae compared to the A1 larvae ( $4.26 \pm 0.12$ ,  $4.29 \pm 0.11$  and  $3.98 \pm 0.09$  mm respectively), while there was no significant difference in eye diameter with an average of  $0.38 \pm 0.08$  mm (Fig. 3). At the end of the second trial, there were no significant differences in standard length and eye diameter between treatments (Fig. 3).

FA composition of eggs, enrichment formulations, enriched rotifers and larvae at the end of the trial are presented in Table 4 for the first trial and Table 5 for the second trial. In addition, the PUFAs content and ratios of the enrichment formulations, enriched rotifers and larvae at the end of the trial are represented in Fig. 4 for trial 1 and Fig. 5 for trial 2.


In the first trial, there was no significant difference in total PUFA between the different enrichments with an average of  $59.1 \pm 1\%$  of total FA. However, total PUFA differed in the rotifers with the greatest amount in A1 rotifers ( $66.2 \pm 1.0\%$  of total FA) followed by Fa rotifers ( $57.6 \pm 1.0\%$  of total FA) and Fb rotifers ( $51.5 \pm 0.2\%$  of total FA). At the end of the trial, total PUFA in the larvae was

**Table 2** Fatty acid profile of fertilized eggs and unfed snook larvae up to 5 days post hatch (DPH). Only fatty acids contributing to at least 1% at one point are reported, all fatty acids are included in totals and ratios. Values are  $\pm$ SEM,  $n = 3$ . Letters indicate significant differences within a same row (Tukey test,  $P < 0.05$ )

% total FA	Fertilized egg	0 DPH	1 DPH	2 DPH	3 DPH	4 DPH	5 DPH
14:0	1.9 $\pm$ 0.1 <sup>bc</sup>	2.4 $\pm$ 0.0 <sup>c</sup>	1.9 $\pm$ 0.1 <sup>bc</sup>	1.9 $\pm$ 0.2 <sup>bc</sup>	0.9 $\pm$ 0.2 <sup>a</sup>	1.3 $\pm$ 0.1 <sup>a</sup>	1.6 $\pm$ 0.1 <sup>b</sup>
16:0	21.7 $\pm$ 0.7	19.7 $\pm$ 0.2	19.4 $\pm$ 0.1	19.9 $\pm$ 0.3	20.2 $\pm$ 0.5	20.3 $\pm$ 0.2	20.6 $\pm$ 0.4
17:0	1.0 $\pm$ 0.0	0.9 $\pm$ 0.0	0.9 $\pm$ 0.0	0.9 $\pm$ 0.0	0.9 $\pm$ 0.0	0.9 $\pm$ 0.1	0.9 $\pm$ 0.1
18:0	5.2 $\pm$ 0.1 <sup>a</sup>	4.5 $\pm$ 0.1 <sup>a</sup>	5.3 $\pm$ 0.3 <sup>a</sup>	6.8 $\pm$ 0.5 <sup>a</sup>	9.2 $\pm$ 0.7 <sup>b</sup>	9.1 $\pm$ 0.1 <sup>b</sup>	8.5 $\pm$ 0.1 <sup>b</sup>
Total saturated	30.6 $\pm$ 0.6 <sup>b</sup>	28.4 $\pm$ 0.4 <sup>a</sup>	28.4 $\pm$ 0.3 <sup>a</sup>	30.3 $\pm$ 0.3 <sup>b</sup>	31.8 $\pm$ 0.4 <sup>bc</sup>	32.5 $\pm$ 0.8 <sup>bc</sup>	32.4 $\pm$ 0.4 <sup>c</sup>
16:1n-7	7.6 $\pm$ 0.3 <sup>b</sup>	9.0 $\pm$ 0.1 <sup>b</sup>	7.4 $\pm$ 0.5 <sup>b</sup>	5.6 $\pm$ 0.4 <sup>b</sup>	3.0 $\pm$ 0.4 <sup>a</sup>	3.8 $\pm$ 0.2 <sup>a</sup>	4.1 $\pm$ 0.4 <sup>a</sup>
18:1n-9	16.9 $\pm$ 0.3 <sup>b</sup>	15.1 $\pm$ 0.2 <sup>b</sup>	13.9 $\pm$ 0.2 <sup>b</sup>	11.8 $\pm$ 0.3 <sup>b</sup>	9.2 $\pm$ 0.3 <sup>a</sup>	9.4 $\pm$ 1.0 <sup>a</sup>	9.7 $\pm$ 1.4 <sup>a</sup>
18:1n-7	4.6 $\pm$ 0.2 <sup>b</sup>	4.3 $\pm$ 0.1 <sup>b</sup>	3.9 $\pm$ 0.1 <sup>b</sup>	3.0 $\pm$ 0.1 <sup>b</sup>	2.4 $\pm$ 0.1 <sup>a</sup>	2.5 $\pm$ 0.2 <sup>a</sup>	2.6 $\pm$ 0.3 <sup>a</sup>
Total monounsaturated	29.5 $\pm$ 0.2 <sup>b</sup>	28.9 $\pm$ 0.2 <sup>b</sup>	25.7 $\pm$ 0.1 <sup>b</sup>	20.9 $\pm$ 0.6 <sup>b</sup>	15.0 $\pm$ 0.8 <sup>a</sup>	16.3 $\pm$ 0.8 <sup>a</sup>	17.1 $\pm$ 0.8 <sup>a</sup>
16:3n-4	1.4 $\pm$ 0.0 <sup>b</sup>	1.4 $\pm$ 0.0 <sup>b</sup>	1.4 $\pm$ 0.0 <sup>b</sup>	1.1 $\pm$ 0.0 <sup>a</sup>	1.1 $\pm$ 0.1 <sup>a</sup>	1.0 $\pm$ 0.6 <sup>a</sup>	1.0 $\pm$ 0.1 <sup>a</sup>
18:2n-6	2.2 $\pm$ 0.5 <sup>b</sup>	2.7 $\pm$ 0.2 <sup>c</sup>	2.7 $\pm$ 0.0 <sup>c</sup>	2.0 $\pm$ 0.0 <sup>b</sup>	1.4 $\pm$ 0.1 <sup>a</sup>	2.1 $\pm$ 0.2 <sup>b</sup>	2.1 $\pm$ 0.8 <sup>b</sup>
20:4n-6	5.4 $\pm$ 0.3 <sup>a</sup>	6.5 $\pm$ 0.2 <sup>ab</sup>	7.7 $\pm$ 0.1 <sup>b</sup>	8.4 $\pm$ 0.1 <sup>b</sup>	9.5 $\pm$ 0.1 <sup>c</sup>	9.5 $\pm$ 0.1 <sup>c</sup>	9.4 $\pm$ 0.2 <sup>c</sup>
20:5n-3	2.4 $\pm$ 0.4 <sup>b</sup>	3.0 $\pm$ 0.1 <sup>a</sup>	2.3 $\pm$ 0.2 <sup>b</sup>	2.3 $\pm$ 0.1 <sup>b</sup>	2.4 $\pm$ 0.1 <sup>b</sup>	2.4 $\pm$ 0.0 <sup>b</sup>	2.3 $\pm$ 0.0 <sup>b</sup>
22:5n-6	2.1 $\pm$ 0.2 <sup>ab</sup>	1.8 $\pm$ 0.0 <sup>a</sup>	2.2 $\pm$ 0.1 <sup>ab</sup>	2.8 $\pm$ 0.2 <sup>ab</sup>	3.0 $\pm$ 0.3 <sup>bc</sup>	3.2 $\pm$ 0.1 <sup>bc</sup>	3.3 $\pm$ 0.2 <sup>c</sup>
22:5n-3	2.7 $\pm$ 0.1 <sup>c</sup>	2.6 $\pm$ 0.0 <sup>c</sup>	2.4 $\pm$ 0.0 <sup>b</sup>	2.0 $\pm$ 0.1 <sup>a</sup>	2.0 $\pm$ 0.1 <sup>a</sup>	2.1 $\pm$ 0.3 <sup>ab</sup>	2.2 $\pm$ 0.7 <sup>b</sup>
22:6n-3	14.5 $\pm$ 0.2 <sup>a</sup>	16.2 $\pm$ 0.2 <sup>b</sup>	17.5 $\pm$ 0.2 <sup>b</sup>	20.0 $\pm$ 0.3 <sup>b</sup>	26.1 $\pm$ 0.6 <sup>c</sup>	26.0 $\pm$ 0.3 <sup>c</sup>	25.8 $\pm$ 0.5 <sup>c</sup>
Total polyunsaturated	33.6 $\pm$ 0.5 <sup>a</sup>	37.4 $\pm$ 0.4 <sup>b</sup>	39.3 $\pm$ 0.4 <sup>b</sup>	41.4 $\pm$ 0.2 <sup>b</sup>	47.7 $\pm$ 0.4 <sup>c</sup>	49.4 $\pm$ 0.9 <sup>c</sup>	48.7 $\pm$ 0.7 <sup>c</sup>
Total n-3	20.8 $\pm$ 0.6 <sup>a</sup>	23.3 $\pm$ 0.3 <sup>a</sup>	23.3 $\pm$ 0.4 <sup>a</sup>	25.3 $\pm$ 0.4 <sup>a</sup>	31.1 $\pm$ 0.6 <sup>b</sup>	31.4 $\pm$ 0.8 <sup>b</sup>	31.4 $\pm$ 0.9 <sup>b</sup>
Total n-6	10.6 $\pm$ 0.5 <sup>a</sup>	11.8 $\pm$ 0.3 <sup>ab</sup>	13.7 $\pm$ 0.2 <sup>ab</sup>	14.1 $\pm$ 0.3 <sup>ab</sup>	14.7 $\pm$ 0.4 <sup>b</sup>	15.5 $\pm$ 0.4 <sup>b</sup>	15.6 $\pm$ 0.9 <sup>b</sup>
n-3/ n-6 ratio	2.1 $\pm$ 0.3 <sup>b</sup>	2.0 $\pm$ 0.1 <sup>ab</sup>	1.7 $\pm$ 0.1 <sup>a</sup>	1.8 $\pm$ 0.1 <sup>a</sup>	2.1 $\pm$ 0.1 <sup>b</sup>	2.0 $\pm$ 0.0 <sup>ab</sup>	2.0 $\pm$ 0.1 <sup>ab</sup>
ARA/EPA ratio	2.8 $\pm$ 0.6 <sup>b</sup>	2.2 $\pm$ 0.1 <sup>a</sup>	3.4 $\pm$ 0.3 <sup>c</sup>	3.6 $\pm$ 0.2 <sup>c</sup>	3.9 $\pm$ 0.2 <sup>d</sup>	4.0 $\pm$ 0.1 <sup>d</sup>	4.0 $\pm$ 0.1 <sup>d</sup>
DHA/EPA ratio	7.1 $\pm$ 0.7 <sup>ab</sup>	5.4 $\pm$ 0.2 <sup>a</sup>	7.7 $\pm$ 0.6 <sup>ab</sup>	8.6 $\pm$ 0.2 <sup>ab</sup>	10.8 $\pm$ 0.1 <sup>b</sup>	10.9 $\pm$ 0.0 <sup>b</sup>	11.1 $\pm$ 0.1 <sup>b</sup>
Total FA (mg FA g <sup>-1</sup> dry weight)	193.3 $\pm$ 3.5 <sup>e</sup>	187.3 $\pm$ 4.1 <sup>de</sup>	163.4 $\pm$ 2.2 <sup>d</sup>	147.7 $\pm$ 1.9 <sup>cd</sup>	134.1 $\pm$ 1.2 <sup>bc</sup>	119.6 $\pm$ 2.3 <sup>ab</sup>	110.1 $\pm$ 2.2 <sup>a</sup>



**Figure 1** Evolution of selected fatty acids in eggs (0 days post fertilization) and unfed pompano larvae (1–5 day post fertilization). Values are  $\pm$ SEM,  $n = 3$ .



**Figure 2** Evolution of selected fatty acids in eggs (0 days post fertilization) and unfed snook larvae (1–5 day post fertilization). Values are  $\pm$ SEM,  $n = 3$ .

significantly higher for A1 and Fa larvae with an average of  $47.4 \pm 0.5\%$  of total FA compared to Fb larvae ( $43.1 \pm 0.7\%$  of total FA). In trial 2, no significant differences were observed in PUFA contents between treatments with an average of  $63.6 \pm 1.0\%$  of total FA in the enrichment,  $54.0 \pm 0.5\%$  of total FA in the rotifers and  $43.5 \pm 0.4\%$  of total FA in the larvae.

DHA content in trial 1 was greatest in the A1 enrichment and rotifers ( $40.8 \pm 0.8\%$  and  $38.2 \pm 0.8\%$  of total FA), followed by the Fa enrichment and rotifers ( $29.8 \pm 0.7\%$  and  $25.0 \pm 0.6\%$  of total FA) then the Fb enrichment and rotifers ( $28.0 \pm 0.4\%$  and  $20.6 \pm 0.3\%$  of total FA). How-

ever, at the end of the trial there was no significant difference in larvae DHA content with an average of  $24.3 \pm 0.4\%$  of total FA. In trial 2, DHA content was significantly higher for the A2 enrichment, rotifers and larvae ( $40.8 \pm 0.6\%$ ,  $27.3 \pm 0.4\%$  and  $22.7 \pm 0.1\%$  of total FA respectively), than in their equivalent from the Fc treatment ( $22.9 \pm 0.2\%$ ,  $15.3 \pm 0.4\%$  and  $19.4 \pm 0.3\%$  of total FA respectively).

ARA content in trial 1 was the greatest in the Fb enrichment ( $17.5 \pm 0.1\%$  of total FA) then in the Fa enrichment ( $14.7 \pm 0.1\%$  of total FA) followed by the A1 enrichment ( $1.6 \pm 0.3\%$  of total FA). Likewise, highest ARA content was observed

**Table 3** Per cent survival and per cent of functional swim bladder at the end of live feed trials with snook larvae fed rotifers enriched with Algamac 3050 (A1), Formulation a (Fa) or Formulation b (Fb) during the first trial, and Algamac 3050 (A2) or Formulation c (Fc) during the second trial. Values are  $\pm$ SEM,  $n = 4$  in trial 1 and trial 2, 10 larvae per tank. Superscript letters indicate significant differences within a same column (Tukey test,  $P < 0.05$ )

		Survival(%)	Functional swim bladder(%)
Trial 1	A1	6.9 $\pm$ 0.9 <sup>b</sup>	80.2 $\pm$ 2.1 <sup>b</sup>
	Fa	2.4 $\pm$ 0.5 <sup>a</sup>	50.9 $\pm$ 2.8 <sup>a</sup>
	Fb	2.1 $\pm$ 0.4 <sup>a</sup>	52.1 $\pm$ 3.4 <sup>a</sup>
Trial 2	A2	5.2 $\pm$ 1.2 <sup>b</sup>	77.1 $\pm$ 3.1 <sup>b</sup>
	Fc	5.4 $\pm$ 0.9 <sup>b</sup>	58.4 $\pm$ 5.2 <sup>a</sup>

in the Fb rotifers (11.1  $\pm$  0.6% of total FA) then in the Fa rotifers (9.3  $\pm$  0.3% of total FA) followed by the A1 rotifers (2.8  $\pm$  0.1% of total FA). At the end of the trial, the ARA content was significantly lower in the A1 larvae (4.4  $\pm$  0.9% of total FA) than in the other treatments (average of 7.1  $\pm$  0.4% of total FA). In trial 2, ARA content was significantly higher in the Fc enrichment, rotifers and larvae (20.8  $\pm$  0.2%, 14.0  $\pm$  0.7% and 8.7  $\pm$  0.3% of total FA respectively) than in their equivalent from the A2 treatment (1.6  $\pm$  0.3%, 3.3  $\pm$  0.2% and 5.2  $\pm$  0.2% of total FA respectively).

At the end of the first trial, the highest DHA/EPA ratio was observed in the A1 larvae (8.9  $\pm$  0.1) and the lowest ratio was observed in the Fa larvae (7.0  $\pm$  0.3) while the Fb ratio was not significantly different from the other treatments (7.9  $\pm$  0.6). In the second trial, the DHA/EPA ratio in the A2 larvae (6.5  $\pm$  0.1) was significantly higher than that of the Fc larvae (5.1  $\pm$  0.3). The highest ARA/EPA ratio in trial 1 was observed in the Fb larvae (2.4  $\pm$  0.1), the lowest ratio in the A1 larvae (1.6  $\pm$  0.2) while the ratio in the Fa larvae was not significantly different from the other treatments (2.0  $\pm$  0.1). In the second trial, the Fb larvae presented an ARA/EPA ratio significantly higher (2.3  $\pm$  0.2) than that of the A2 larvae (1.5  $\pm$  0.2).


## Discussion

The study of FA utilization in pompano and snook unfed larvae revealed clear patterns of lipid utilization and mobilization. Eggs from both species contained an oil globule and are classified as being high in lipids (>19% of dry weight) (Sargent, Henderson & Tocher 1989). After hatching the decline in total FA reflected the utilization of lipids

as an energy source. This is in accordance with previous studies in species such as red drum *Sciaenops ocellatus* (Vetter, Hodson & Arnold 1983), red sea bream *Pagrus major* (Tandler, Watanabe, Satoh & Fukusho 1989), gilthead sea bream (Koven, Kissil & Tandler 1989; Rønnestad, Koven, Tandler, Harel & Fyhn 1994), turbot (Rainuzzo, Reitan, Jørgensen, Olsen & Jørgensen 1994), Senegal sole (Mourente & Vázquez 1996), common dentex *Dentex dentex* (Mourente, Rodriguez, Grau & Pastor 1999), white sea bream *Diplodus sargus* (Cejas, Almansa, Jérez, Bolaños, Felipe & Lorenzo 2004) and Atlantic bluefin tuna *Thunnus thynnus* (Morais, Mourente, Ortega, Tocher & Tocher 2011) indicating that fast developing eggs from temperate and warm waters, with high lipid content and an oil globule, incorporate a large fraction of neutral lipids in the oil droplet which is utilized for energy soon after hatching. Contrastingly, cold water species with slower egg development such as Atlantic herring *Clupea harengus* (Tocher, Fraser, Sargent & Gamble 1985a) Atlantic cod *Gadus morhua* and plaice *Pleuronectes platessa* (Rainuzzo, Reitan & Jørgensen 1992) or Atlantic halibut *Hippoglossus hippoglossus* (Rainuzzo *et al.* 1992; Rønnestad, Finn, Lein & Lie 1995) mainly catabolize free amino acid after hatching (Rønnestad 1999).

The neutral lipid fraction is generally rich in mono-unsaturated FAs that are preferentially used for energy (Kamler 2007). It seems to be the case in this study as, even though no lipid class analyses were performed, a strong decrease in mono-unsaturated FAs and no increase in their elongation products were observed in both species after hatching. The selective retention of certain fatty acids has been described as a biochemical strategy allowing the preservation of the most essential components of biological membranes during starvation periods (Izquierdo 1996). It has been recently demonstrated in turbot that the expression profile of some genes involved in lipid metabolism (Hepatic lipase, FA synthetase and Diacylglycerol O-acyltransferase homologue 1) responded solely to starvation (Cunha, Galante-Oliveira, Rocha, Planas, Urbatzka & Castro 2013). In the present study saturated fatty acids (SFAs) and PUFAs were preferentially retained. SFAs and PUFAs play important roles in respectively the sn-1 and sn-2 position of structural phospholipids with phosphatidylcholine containing high levels of 16:0, phosphatidylserine incorporating high levels of 18:0 associated with C22 PUFAs,





**Figure 3** Standard length (A) and eye diameter (B) of snook larvae fed rotifers enriched with Algamac 3050 (A1), Formulation a (Fa1) or Formulation b (Fb1) during the first trial, and Algamac 3050 (A2) or Formulation c (Fc2) during the second trial. Values are  $\pm$ SE of the mean,  $n = 4$  in trial 1 and trial 2, 10 larvae per tank and sampling point. Letters indicate significant differences between treatments and time points (Tukey test,  $P < 0.05$ ).

and phosphatidylinositol including high levels of 18:0 associated with C20 PUFAs and more particularly ARA (Tocher 1995; Sargent *et al.* 2002). In both pompano and snook, the main PUFAs selectively retained were DHA and ARA, while low levels of EPA were observed. DHA is selectively retained in unfed larvae of several species, including Atlantic herring (Tocher, Fraser, Sargent & Gamble 1985b), gilthead sea bream (Koven *et al.* 1989), red sea bream (Tandler *et al.* 1989), cod (Van der Meer, Wilhelmsen, Klungsosyr & Kvenseth 1993), turbot (Rainuzzo *et al.* 1994), Senegal sole (Mourente & Vázquez 1996) common dentex (Mourente *et al.* 1999) and Atlantic bluefin tuna (Morais *et al.* 2011). This reflects the importance of DHA in larval structural development and in particular on


neural and visual functions (Bell, Castell *et al.* 1995; Bell, Batty *et al.* 1995; Sargent, Bell *et al.* 1999). ARA is selectively retained along with DHA in fewer species, including turbot (Rainuzzo *et al.* 1994), Senegal sole (Mourente & Vázquez 1996) and bluefin tuna (Morais *et al.* 2011). In addition, in Atlantic halibut, cod, plaice (Rainuzzo *et al.* 1992) and white sea bream (Cejas *et al.* 2004), ARA solely was conserved, while utilization of DHA was observed, indicating a special role of ARA in the early larval development of these species. Concurrently to the retention of ARA, a decrease in EPA level is observed throughout the starvation period in pompano while a significant decrease is only observed between 0 and 1 dph in snook. In the absence of information on bioconversion rates in

**Table 4** Fatty acid profile of fertilized eggs, Algamac 3050 (A1), Formulation a (Fa) and Formulation b (Fb) and their enriched rotifers during the trial (6 sampling times) as well as their resulting snook larvae at the end of the trial. Only fatty acids contributing to at least 1% in one of the treatments are reported, all fatty acids are included in totals and ratios. Values are  $\pm$ SEM,  $n = 3$  for the eggs and enrichments,  $n = 4$  for larvae (50 larvae per tank),  $n = 6$  for enriched rotifers. Superscripted letters indicate significant differences within a same row (Tukey test,  $P < 0.05$ ). nd means not detected

% total FA	A1			Fa			Fb			
	Fertilized Eggs	Enrichment	Enriched Rotifers	Larvae	Enrichment	Enriched Rotifers	Larvae	Enrichment	Enriched Rotifers	Larvae
12:0	0.0 $\pm$ 0.0 <sup>a</sup>	0.3 $\pm$ 0.1 <sup>b</sup>	0.1 $\pm$ 0.0a <sup>b</sup>	0.2 $\pm$ 0.1 <sup>b</sup>	0.9 $\pm$ 0.0 <sup>c</sup>	0.2 $\pm$ 0.0 <sup>b</sup>	0.2 $\pm$ 0.0 <sup>b</sup>	1.1 $\pm$ 0.0 <sup>c</sup>	0.3 $\pm$ 0.0 <sup>b</sup>	0.3 $\pm$ 0.1 <sup>b</sup>
14:0	3.1 $\pm$ 0.1 <sup>b</sup>	9.8 $\pm$ 0.6 <sup>e</sup>	4.4 $\pm$ 0.2 <sup>c</sup>	1.3 $\pm$ 0.1 <sup>a</sup>	6.0 $\pm$ 0.2 <sup>d</sup>	3.2 $\pm$ 0.1 <sup>b</sup>	0.9 $\pm$ 0.2 <sup>a</sup>	6.0 $\pm$ 0.1 <sup>d</sup>	3.4 $\pm$ 0.2 <sup>bc</sup>	0.9 $\pm$ 0.1 <sup>a</sup>
16:0	17.9 $\pm$ 0.8 <sup>c</sup>	25.9 $\pm$ 0.7 <sup>d</sup>	15.2 $\pm$ 0.8 <sup>bc</sup>	16.5 $\pm$ 1.2 <sup>c</sup>	11.7 $\pm$ 0.3 <sup>ab</sup>	12.0 $\pm$ 0.3 <sup>ab</sup>	14.6 $\pm$ 0.3 <sup>bc</sup>	11.8 $\pm$ 0.2 <sup>ab</sup>	11.0 $\pm$ 0.5 <sup>a</sup>	16.2 $\pm$ 0.6 <sup>c</sup>
17:0	0.7 $\pm$ 0.0 <sup>cd</sup>	nd	0.5 $\pm$ 0.0 <sup>c</sup>	1.2 $\pm$ 0.2 <sup>ef</sup>	0.1 $\pm$ 0.0 <sup>b</sup>	0.9 $\pm$ 0.0 <sup>de</sup>	1.2 $\pm$ 0.1 <sup>ef</sup>	0.1 $\pm$ 0.0 <sup>b</sup>	0.7 $\pm$ 0.1 <sup>cd</sup>	1.3 $\pm$ 0.1 <sup>f</sup>
18:0	4.8 $\pm$ 0.1 <sup>c</sup>	0.5 $\pm$ 0.0 <sup>a</sup>	1.5 $\pm$ 0.1 <sup>b</sup>	8.4 $\pm$ 0.4 <sup>d</sup>	1.7 $\pm$ 0.6 <sup>b</sup>	2.7 $\pm$ 0.1 <sup>b</sup>	7.9 $\pm$ 0.5 <sup>d</sup>	2.2 $\pm$ 0.6 <sup>b</sup>	2.6 $\pm$ 0.2 <sup>b</sup>	9.3 $\pm$ 0.7 <sup>d</sup>
Total saturated	27.1 $\pm$ 0.8 <sup>c</sup>	37.0 $\pm$ 0.6 <sup>d</sup>	22.0 $\pm$ 0.8 <sup>ab</sup>	27.9 $\pm$ 0.7 <sup>c</sup>	20.4 $\pm$ 0.2 <sup>a</sup>	19.2 $\pm$ 0.4 <sup>a</sup>	25.0 $\pm$ 0.7 <sup>bc</sup>	21.3 $\pm$ 0.9 <sup>ab</sup>	18.2 $\pm$ 0.8 <sup>a</sup>	28.4 $\pm$ 1.5 <sup>c</sup>
16:1n-7	8.0 $\pm$ 0.2 <sup>f</sup>	nd	3.2 $\pm$ 0.2 <sup>c</sup>	3.0 $\pm$ 0.2 <sup>c</sup>	2.2 $\pm$ 0.0 <sup>b</sup>	5.9 $\pm$ 0.2 <sup>e</sup>	3.2 $\pm$ 0.4 <sup>c</sup>	1.7 $\pm$ 0.0 <sup>b</sup>	4.8 $\pm$ 0.4 <sup>d</sup>	2.9 $\pm$ 0.5 <sup>c</sup>
18:1n-9	11.1 $\pm$ 0.4 <sup>f</sup>	0.1 $\pm$ 0.0 <sup>a</sup>	0.6 $\pm$ 0.1 <sup>b</sup>	4.9 $\pm$ 0.3 <sup>c</sup>	14.5 $\pm$ 0.2 <sup>g</sup>	7.7 $\pm$ 0.3 <sup>de</sup>	6.9 $\pm$ 0.3 <sup>d</sup>	13.9 $\pm$ 0.3 <sup>g</sup>	8.4 $\pm$ 0.3 <sup>e</sup>	7.1 $\pm$ 0.3 <sup>de</sup>
18:1n-7	6.6 $\pm$ 0.1 <sup>d</sup>	nd	1.8 $\pm$ 0.1 <sup>b</sup>	3.2 $\pm$ 0.2 <sup>c</sup>	nd	3.8 $\pm$ 0.2 <sup>=</sup>	3.7 $\pm$ 0.1 <sup>c</sup>	nd	3.1 $\pm$ 0.3 <sup>c</sup>	2.9 $\pm$ 0.3 <sup>bc</sup>
20:1n-9	0.5 $\pm$ 0.0 <sup>bc</sup>	nd	0.5 $\pm$ 0.1 <sup>bc</sup>	0.7 $\pm$ 0.1 <sup>c</sup>	0.3 $\pm$ 0.0 <sup>bc</sup>	1.4 $\pm$ 0.2 <sup>d</sup>	1.4 $\pm$ 0.1 <sup>d</sup>	0.2 $\pm$ 0.0 <sup>b</sup>	1.4 $\pm$ 0.1 <sup>d</sup>	1.3 $\pm$ 0.1 <sup>d</sup>
Total monounsaturated	26.4 $\pm$ 0.5 <sup>f</sup>	0.1 $\pm$ 0.0 <sup>a</sup>	6.2 $\pm$ 0.4 <sup>b</sup>	11.9 $\pm$ 0.7 <sup>c</sup>	17.0 $\pm$ 0.3 <sup>de</sup>	18.9 $\pm$ 0.4 <sup>e</sup>	15.3 $\pm$ 0.4 <sup>d</sup>	15.9 $\pm$ 0.3 <sup>d</sup>	17.8 $\pm$ 0.4 <sup>de</sup>	14.6 $\pm$ 0.6 <sup>d</sup>
18:2n-6	1.8 $\pm$ 0.0 <sup>c</sup>	0.1 $\pm$ 0.0 <sup>a</sup>	0.9 $\pm$ 0.1 <sup>b</sup>	1.5 $\pm$ 0.1 <sup>b</sup>	4.6 $\pm$ 0.1 <sup>e</sup>	4.4 $\pm$ 0.1 <sup>e</sup>	2.4 $\pm$ 0.1 <sup>d</sup>	4.9 $\pm$ 0.1 <sup>e</sup>	4.6 $\pm$ 0.1 <sup>e</sup>	2.6 $\pm$ 0.2 <sup>d</sup>
18:3n-6	0.2 $\pm$ 0.1 <sup>a</sup>	0.2 $\pm$ 0.0 <sup>a</sup>	0.3 $\pm$ 0.0 <sup>a</sup>	0.5 $\pm$ 0.0 <sup>b</sup>	0.8 $\pm$ 0.0 <sup>c</sup>	0.7 $\pm$ 0.0 <sup>c</sup>	0.5 $\pm$ 0.1 <sup>b</sup>	1.1 $\pm$ 0.1 <sup>d</sup>	0.8 $\pm$ 0.0 <sup>c</sup>	0.6 $\pm$ 0.1 <sup>b</sup>
18:4n-3	0.5 $\pm$ 0.1 <sup>a</sup>	0.3 $\pm$ 0.1 <sup>a</sup>	0.4 $\pm$ 0.1 <sup>a</sup>	1.1 $\pm$ 0.8 <sup>ab</sup>	2.4 $\pm$ 0.1 <sup>c</sup>	1.7 $\pm$ 0.2 <sup>bc</sup>	0.4 $\pm$ 0.1 <sup>a</sup>	0.7 $\pm$ 0.0 <sup>ab</sup>	0.8 $\pm$ 0.1 <sup>ab</sup>	0.4 $\pm$ 0.1 <sup>a</sup>
20:3n-6	0.6 $\pm$ 0.1 <sup>bc</sup>	0.4 $\pm$ 0.0 <sup>a</sup>	0.8 $\pm$ 0.1 <sup>c</sup>	0.8 $\pm$ 0.1 <sup>c</sup>	0.5 $\pm$ 0.2 <sup>b</sup>	1.6 $\pm$ 0.1 <sup>e</sup>	1.0 $\pm$ 0.1 <sup>d</sup>	1.1 $\pm$ 0.2 <sup>d</sup>	1.7 $\pm$ 0.2 <sup>e</sup>	0.8 $\pm$ 0.1 <sup>c</sup>
20:4n-6	5.1 $\pm$ 0.0 <sup>c</sup>	1.6 $\pm$ 0.3 <sup>a</sup>	2.8 $\pm$ 0.1 <sup>b</sup>	4.4 $\pm$ 0.9 <sup>c</sup>	14.7 $\pm$ 0.1 <sup>g</sup>	9.3 $\pm$ 0.3 <sup>e</sup>	7.1 $\pm$ 0.2 <sup>d</sup>	17.5 $\pm$ 0.1 <sup>h</sup>	11.1 $\pm$ 0.6 <sup>f</sup>	7.0 $\pm$ 0.6 <sup>d</sup>
20:4n-3	0.4 $\pm$ 0.0 <sup>bc</sup>	0.8 $\pm$ 0.0 <sup>cd</sup>	1.0 $\pm$ 0.1 <sup>d</sup>	0.4 $\pm$ 0.1 <sup>bc</sup>	nd	0.7 $\pm$ 0.1 <sup>cd</sup>	0.3 $\pm$ 0.1 <sup>bc</sup>	nd	0.5 $\pm$ 0.0 <sup>c</sup>	0.2 $\pm$ 0.1 <sup>b</sup>
20:5n-3	4.5 $\pm$ 0.1 <sup>c</sup>	1.0 $\pm$ 0.5 <sup>a</sup>	5.8 $\pm$ 0.5 <sup>d</sup>	2.8 $\pm$ 0.3 <sup>b</sup>	1.0 $\pm$ 0.0 <sup>=</sup>	7.7 $\pm$ 0.4 <sup>e</sup>	3.6 $\pm$ 0.3 <sup>bc</sup>	1.2 $\pm$ 0.0 <sup>a</sup>	6.4 $\pm$ 0.2 <sup>d</sup>	3.0 $\pm$ 0.3 <sup>b</sup>
22:5n-6	1.5 $\pm$ 0.1 <sup>c</sup>	14.6 $\pm$ 0.5 <sup>e</sup>	12.8 $\pm$ 0.3 <sup>e</sup>	8.8 $\pm$ 0.2 <sup>d</sup>	0.2 $\pm$ 0.0 <sup>ab</sup>	0.6 $\pm$ 0.1 <sup>b</sup>	1.1 $\pm$ 0.4 <sup>bc</sup>	0.1 $\pm$ 0.0 <sup>a</sup>	0.5 $\pm$ 0.1 <sup>b</sup>	1.3 $\pm$ 0.9 <sup>bc</sup>
22:5n-3	3.3 $\pm$ 0.1 <sup>cd</sup>	0.3 $\pm$ 0.0 <sup>a</sup>	2.7 $\pm$ 0.3 <sup>c</sup>	1.9 $\pm$ 0.1 <sup>b</sup>	2.5 $\pm$ 0.1 <sup>bc</sup>	4.4 $\pm$ 0.2 <sup>d</sup>	3.2 $\pm$ 0.3 <sup>cd</sup>	3.0 $\pm$ 0.0 <sup>cd</sup>	3.5 $\pm$ 0.4 <sup>cd</sup>	2.3 $\pm$ 0.4 <sup>bc</sup>
22:6n-3	22.3 $\pm$ 0.8 <sup>ab</sup>	40.8 $\pm$ 0.8 <sup>e</sup>	38.2 $\pm$ 0.8 <sup>e</sup>	24.5 $\pm$ 0.4 <sup>b</sup>	29.8 $\pm$ 0.7 <sup>d</sup>	25.0 $\pm$ 1.0 <sup>b</sup>	25.1 $\pm$ 0.6 <sup>b</sup>	28.0 $\pm$ 0.4 <sup>c</sup>	20.6 $\pm$ 0.3 <sup>a</sup>	23.4 $\pm$ 0.3 <sup>ab</sup>
Total polyunsaturated	42.6 $\pm$ 0.9 <sup>a</sup>	60.4 $\pm$ 1.4 <sup>d</sup>	66.2 $\pm$ 1.0 <sup>e</sup>	48.2 $\pm$ 0.7 <sup>b</sup>	58.2 $\pm$ 1.0 <sup>d</sup>	57.6 $\pm$ 1.0 <sup>d</sup>	46.7 $\pm$ 0.4 <sup>b</sup>	58.7 $\pm$ 0.5 <sup>d</sup>	51.5 $\pm$ 0.2 <sup>c</sup>	43.1 $\pm$ 0.7 <sup>a</sup>
Total n-3	31.8 $\pm$ 0.6 <sup>b</sup>	43.4 $\pm$ 0.6 <sup>cd</sup>	48.4 $\pm$ 0.9 <sup>d</sup>	31.2 $\pm$ 3.2 <sup>b</sup>	37.1 $\pm$ 0.8 <sup>c</sup>	35.8 $\pm$ 1.0 <sup>bc</sup>	31.2 $\pm$ 1.2 <sup>b</sup>	33.8 $\pm$ 0.5 <sup>b</sup>	36.8 $\pm$ 0.6 <sup>c</sup>	26.4 $\pm$ 1.5 <sup>a</sup>
Total n-6	9.4 $\pm$ 0.1 <sup>a</sup>	16.8 $\pm$ 0.4 <sup>c</sup>	17.7 $\pm$ 0.3 <sup>cd</sup>	16.2 $\pm$ 1.1 <sup>c</sup>	21.0 $\pm$ 0.2 <sup>d</sup>	17.0 $\pm$ 0.4 <sup>c</sup>	12.6 $\pm$ 0.3 <sup>b</sup>	24.9 $\pm$ 0.0 <sup>e</sup>	19.0 $\pm$ 0.5 <sup>cd</sup>	12.5 $\pm$ 1.6 <sup>b</sup>
n-3/n-6	3.4 $\pm$ 0.2 <sup>f</sup>	2.6 $\pm$ 0.1 <sup>e</sup>	2.7 $\pm$ 0.0 <sup>e</sup>	1.9 $\pm$ 0.2 <sup>c</sup>	1.8 $\pm$ 0.0 <sup>b</sup>	2.1 $\pm$ 0.1 <sup>d</sup>	2.5 $\pm$ 0.2 <sup>de</sup>	1.4 $\pm$ 0.0 <sup>a</sup>	1.9 $\pm$ 0.1 <sup>c</sup>	2.1 $\pm$ 0.0 <sup>d</sup>
ARA/EPA	1.1 $\pm$ 0.0 <sup>b</sup>	1.5 $\pm$ 0.2 <sup>c</sup>	0.5 $\pm$ 0.0 <sup>a</sup>	1.6 $\pm$ 0.2 <sup>cd</sup>	14.9 $\pm$ 0.1 <sup>f</sup>	1.2 $\pm$ 0.1 <sup>b</sup>	2.0 $\pm$ 0.1 <sup>de</sup>	14.7 $\pm$ 0.6 <sup>f</sup>	1.8 $\pm$ 0.2 <sup>d</sup>	2.4 $\pm$ 0.1 <sup>e</sup>
DHA/EPA	5.0 $\pm$ 0.2 <sup>ab</sup>	39.4 $\pm$ 1.5 <sup>f</sup>	6.6 $\pm$ 0.8 <sup>b</sup>	8.9 $\pm$ 0.1 <sup>c</sup>	30.2 $\pm$ 0.5 <sup>e</sup>	3.3 $\pm$ 0.2 <sup>a</sup>	7.0 $\pm$ 0.3 <sup>b</sup>	23.5 $\pm$ 0.4 <sup>d</sup>	3.2 $\pm$ 0.2 <sup>a</sup>	7.9 $\pm$ 0.6 <sup>bc</sup>
Total FA (mg FA g <sup>-1</sup> dry weight)	151.4 $\pm$ 5.0 <sup>d</sup>	398.3 $\pm$ 5.0 <sup>e</sup>	153.9 $\pm$ 6.3 <sup>d</sup>	88.5 $\pm$ 6.3 <sup>b</sup>	254.8 $\pm$ 1.2 <sup>e</sup>	82.7 $\pm$ 3.9 <sup>b</sup>	64.2 $\pm$ 3.5 <sup>a</sup>	328.7 $\pm$ 4.1 <sup>f</sup>	106.2 $\pm$ 8.4 <sup>c</sup>	94.5 $\pm$ 2.2 <sup>b</sup>

**Table 5** Fatty acid profile of fertilized eggs, Algamac 3050 (A2) and Formulation c (Fc), and their enriched rotifers during the trial (six sampling times) as well as their resulting snook larvae at the end of the trial. Only fatty acids contributing to at least 1% in one of the treatments are reported, all fatty acids are included in totals and ratios. Values are  $\pm$ SEM,  $n = 3$  for the eggs and enrichments,  $n = 4$  for larvae (50 larvae per tank),  $n = 6$  for enriched rotifers. Superscripted letters indicate significant differences within a same row (Tukey test,  $P < 0.05$ ). nd means not detected

% total FA	A2			Fc			
	Fertilized Eggs	Enrichment	Enriched Rotifers	Larvae	Enrichment	Enriched Rotifers	Larvae
14:0	2.2 $\pm$ 0.1 <sup>b</sup>	9.8 $\pm$ 0.6 <sup>d</sup>	5.1 $\pm$ 0.1 <sup>c</sup>	1.2 $\pm$ 0.0 <sup>a</sup>	5.8 $\pm$ 0.3 <sup>c</sup>	3.0 $\pm$ 0.1 <sup>b</sup>	1.2 $\pm$ 0.1 <sup>a</sup>
16:0	22.9 $\pm$ 0.5 <sup>b</sup>	25.9 $\pm$ 0.7 <sup>c</sup>	18.0 $\pm$ 0.3 <sup>a</sup>	23.8 $\pm$ 0.1 <sup>bc</sup>	16.0 $\pm$ 0.4 <sup>a</sup>	15.8 $\pm$ 0.3 <sup>a</sup>	22.3 $\pm$ 0.3 <sup>b</sup>
18:0	5.5 $\pm$ 0.1 <sup>d</sup>	0.5 $\pm$ 0.1 <sup>a</sup>	1.7 $\pm$ 0.0 <sup>b</sup>	8.9 $\pm$ 0.1 <sup>e</sup>	3.0 $\pm$ 0.1 <sup>c</sup>	2.6 $\pm$ 0.1 <sup>c</sup>	8.0 $\pm$ 0.2 <sup>e</sup>
Total saturated	32.4 $\pm$ 0.5 <sup>c</sup>	37.0 $\pm$ 1.1 <sup>d</sup>	25.6 $\pm$ 0.4 <sup>b</sup>	34.9 $\pm$ 0.6 <sup>c</sup>	25.5 $\pm$ 0.5 <sup>b</sup>	21.9 $\pm$ 0.3 <sup>a</sup>	32.5 $\pm$ 0.3 <sup>c</sup>
16:1n7	8.0 $\pm$ 0.4 <sup>c</sup>	nd	7.7 $\pm$ 0.2 <sup>c</sup>	4.0 $\pm$ 0.1 <sup>b</sup>	0.6 $\pm$ 0.1 <sup>a</sup>	8.2 $\pm$ 0.2 <sup>c</sup>	5.1 $\pm$ 0.2 <sup>bc</sup>
18:1n9	14.4 $\pm$ 0.8 <sup>e</sup>	0.1 $\pm$ 0.0 <sup>a</sup>	2.7 $\pm$ 0.1 <sup>b</sup>	5.6 $\pm$ 0.2 <sup>b</sup>	2.0 $\pm$ 1.1 <sup>b</sup>	6.3 $\pm$ 0.2 <sup>c</sup>	7.1 $\pm$ 0.3 <sup>d</sup>
18:1n7	5.3 $\pm$ 0.1 <sup>d</sup>	nd	2.3 $\pm$ 0.1 <sup>b</sup>	2.9 $\pm$ 0.2 <sup>b</sup>	3.9 $\pm$ 1.1 <sup>c</sup>	2.3 $\pm$ 0.1 <sup>b</sup>	2.8 $\pm$ 0.1 <sup>b</sup>
20:1n9	0.2 $\pm$ 0.0 <sup>b</sup>	nd	1.2 $\pm$ 0.0 <sup>c</sup>	1.0 $\pm$ 0.0 <sup>c</sup>	0.2 $\pm$ 0.0 <sup>b</sup>	1.4 $\pm$ 0.0 <sup>c</sup>	1.2 $\pm$ 0.1 <sup>c</sup>
Total monounsaturated	28.2 $\pm$ 0.6 <sup>e</sup>	0.1 $\pm$ 0.0 <sup>a</sup>	13.9 $\pm$ 0.3 <sup>c</sup>	13.6 $\pm$ 0.1 <sup>c</sup>	6.7 $\pm$ 0.3 <sup>b</sup>	18.2 $\pm$ 0.3 <sup>d</sup>	16.3 $\pm$ 0.3 <sup>d</sup>
16:3n4	1.5 $\pm$ 0.1 <sup>d</sup>	0.2 $\pm$ 0.0 <sup>b</sup>	0.2 $\pm$ 0.0 <sup>b</sup>	0.3 $\pm$ 0.0 <sup>c</sup>	nd	0.3 $\pm$ 0.0 <sup>c</sup>	0.3 $\pm$ 0.0 <sup>c</sup>
18:2n6	1.9 $\pm$ 0.1 <sup>bc</sup>	0.1 $\pm$ 0.0 <sup>a</sup>	1.6 $\pm$ 0.1 <sup>b</sup>	1.7 $\pm$ 0.1 <sup>b</sup>	4.7 $\pm$ 0.1 <sup>d</sup>	4.8 $\pm$ 0.1 <sup>d</sup>	2.6 $\pm$ 0.2 <sup>c</sup>
18:3n6	0.6 $\pm$ 0.0 <sup>c</sup>	0.2 $\pm$ 0.0 <sup>a</sup>	0.4 $\pm$ 0.1 <sup>bc</sup>	0.3 $\pm$ 0.0 <sup>ab</sup>	1.1 $\pm$ 0.0 <sup>d</sup>	1.0 $\pm$ 0.0 <sup>d</sup>	0.5 $\pm$ 0.0 <sup>c</sup>
18:3n3	1.1 $\pm$ 0.1 <sup>c</sup>	0.1 $\pm$ 0.1 <sup>ab</sup>	0.2 $\pm$ 0.0 <sup>ab</sup>	0.1 $\pm$ 0.0 <sup>a</sup>	0.2 $\pm$ 0.0 <sup>ab</sup>	0.3 $\pm$ 0.0 <sup>b</sup>	0.1 $\pm$ 0.0 <sup>a</sup>
20:3n6	0.5 $\pm$ 0.0 <sup>b</sup>	0.4 $\pm$ 0.0 <sup>a</sup>	0.7 $\pm$ 0.1 <sup>bc</sup>	0.6 $\pm$ 0.0 <sup>b</sup>	1.8 $\pm$ 0.1 <sup>d</sup>	1.6 $\pm$ 0.0 <sup>d</sup>	0.9 $\pm$ 0.1 <sup>c</sup>
20:4n6	4.7 $\pm$ 0.2 <sup>bc</sup>	1.6 $\pm$ 0.3 <sup>a</sup>	3.3 $\pm$ 0.2 <sup>b</sup>	5.2 $\pm$ 0.2 <sup>c</sup>	20.8 $\pm$ 0.2 <sup>f</sup>	14.0 $\pm$ 0.7 <sup>e</sup>	8.7 $\pm$ 0.3 <sup>d</sup>
20:5n3	2.4 $\pm$ 0.1 <sup>b</sup>	1.0 $\pm$ 0.5 <sup>a</sup>	7.3 $\pm$ 0.3 <sup>d</sup>	3.5 $\pm$ 0.1 <sup>c</sup>	1.4 $\pm$ 0.1 <sup>a</sup>	7.1 $\pm$ 0.4 <sup>d</sup>	3.8 $\pm$ 0.2 <sup>c</sup>
22:5n6	1.3 $\pm$ 0.1 <sup>a</sup>	14.6 $\pm$ 0.5 <sup>e</sup>	8.9 $\pm$ 0.2 <sup>d</sup>	5.5 $\pm$ 0.2 <sup>c</sup>	5.8 $\pm$ 0.2 <sup>c</sup>	3.4 $\pm$ 0.1 <sup>b</sup>	3.1 $\pm$ 0.1 <sup>b</sup>
22:5n3	2.9 $\pm$ 0.1 <sup>b</sup>	0.3 $\pm$ 0.0 <sup>a</sup>	2.5 $\pm$ 0.1 <sup>b</sup>	2.5 $\pm$ 0.1 <sup>b</sup>	4.6 $\pm$ 0.3 <sup>d</sup>	3.7 $\pm$ 0.1 <sup>c</sup>	3.1 $\pm$ 0.0 <sup>bc</sup>
22:6n3	9.5 $\pm$ 0.4 <sup>a</sup>	40.8 $\pm$ 0.6 <sup>f</sup>	27.3 $\pm$ 0.4 <sup>e</sup>	22.7 $\pm$ 0.1 <sup>d</sup>	22.9 $\pm$ 0.2 <sup>d</sup>	15.3 $\pm$ 0.4 <sup>b</sup>	19.4 $\pm$ 0.3 <sup>c</sup>
Total polyunsaturated	29.3 $\pm$ 0.6 <sup>a</sup>	62.4 $\pm$ 1.4 <sup>d</sup>	54.4 $\pm$ 0.4 <sup>c</sup>	43.4 $\pm$ 0.1 <sup>b</sup>	64.77 $\pm$ 0.5 <sup>d</sup>	53.5 $\pm$ 0.6 <sup>c</sup>	43.7 $\pm$ 0.6 <sup>b</sup>
Total $\omega$ 3	17.5 $\pm$ 0.4 <sup>a</sup>	43.4 $\pm$ 1.4 <sup>d</sup>	38.8 $\pm$ 0.3 <sup>d</sup>	29.3 $\pm$ 0.1 <sup>c</sup>	30.26 $\pm$ 0.9 <sup>c</sup>	27.6 $\pm$ 0.2 <sup>bc</sup>	26.9 $\pm$ 0.1 <sup>b</sup>
Total $\omega$ 6	9.3 $\pm$ 0.2 <sup>a</sup>	16.8 $\pm$ 0.4 <sup>c</sup>	15.0 $\pm$ 0.1 <sup>b</sup>	13.4 $\pm$ 0.1 <sup>b</sup>	34.44 $\pm$ 0.3 <sup>e</sup>	25.1 $\pm$ 0.7 <sup>d</sup>	16.1 $\pm$ 0.6 <sup>c</sup>
$\omega$ 3/ $\omega$ 6	1.9 $\pm$ 0.2 <sup>cd</sup>	2.6 $\pm$ 0.1 <sup>e</sup>	2.6 $\pm$ 0.2 <sup>e</sup>	2.2 $\pm$ 0.4 <sup>d</sup>	0.88 $\pm$ 0.0 <sup>a</sup>	1.1 $\pm$ 0.1 <sup>b</sup>	1.7 $\pm$ 0.1 <sup>c</sup>
ARA/EPA	2.0 $\pm$ 0.1 <sup>c</sup>	1.5 $\pm$ 0.2 <sup>b</sup>	0.5 $\pm$ 0.1 <sup>a</sup>	1.5 $\pm$ 0.2 <sup>b</sup>	14.53 $\pm$ 0.3 <sup>d</sup>	2.0 $\pm$ 0.1 <sup>c</sup>	2.3 $\pm$ 0.2 <sup>c</sup>
DHA/EPA	4.0 $\pm$ 0.3 <sup>b</sup>	39.4 $\pm$ 0.5 <sup>f</sup>	3.7 $\pm$ 0.2 <sup>b</sup>	6.5 $\pm$ 0.1 <sup>d</sup>	15.95 $\pm$ 0.4 <sup>e</sup>	2.2 $\pm$ 0.1 <sup>a</sup>	5.1 $\pm$ 0.3 <sup>c</sup>
Total FA (mg FA g <sup>-1</sup> dry weight)	173.4 $\pm$ 2.9 <sup>c</sup>	398.3 $\pm$ 5.0 <sup>d</sup>	174.8 $\pm$ 4.1 <sup>c</sup>	110.0 $\pm$ 5.5 <sup>a</sup>	392.0 $\pm$ 2.0 <sup>d</sup>	141.5 $\pm$ 3.1 <sup>b</sup>	120.3 $\pm$ 2.9 <sup>ab</sup>




**Figure 4** DHA, EPA, ARA, DHA/EPA and ARA/EPA ratios in the enrichment formulations, enriched rotifers and larvae at the end of the first trial. Values are  $\pm$ SEM,  $n = 3$  for the enrichment formulations,  $n = 4$  for larvae (50 larvae per tank) and  $n = 6$  for enriched rotifers. Letters indicate significant differences between treatments.

these species, the decrease could be due to either catabolism or bioconversion to DHA.

Among the enrichment formulations tested in the second part of the study, the ARA supplementation did not increase survival or eye diameter. An increase in growth was observed in the first enrichment trial, yet, considering the lower survival and swim bladder inflation rates in the supplemented treatments, the difference can probably be attributed to the lower density in these treatments rather than the supplementation. In the second trial, no significant difference was observed in growth or survival between treatments even though the rate of functional swim bladder was significantly lower for the supplemented larvae. The oil-based formulation of the supplemented treatments seemed to negatively impact the swim bladder inflation success of the larvae compared to the flake formulation of Al-

gamac 3050 despite efforts to reduce the oil film at the surface of the tanks. Future studies could benefit from supplementing the enrichment with ARA powder rather than oil.

Large differences were observed in the fatty acid profile of the larvae at the end of the trials. The main challenge in achieving a high ARA/EPA ratio is the relatively high natural EPA content of rotifers. Indeed, even when the proportion of EPA in the enrichment formulation was as low as 1%, the proportion reached over 5% in the enriched rotifers. At the end of the first trial, Fa and Fb larvae successfully incorporated ARA and EPA at a 2:1 ratio; however, the low 22:5n6 (Docosapentaenoic acid, DPA) content in these treatments compared to that of the Algamac treatment was a concern. In the second trial, a higher level of DPA was provided in the Fc formulation, which might



**Figure 5** DHA, EPA, ARA, DHA/EPA and ARA/EPA ratios in the enrichment formulations, enriched rotifers and larvae at the end of the second trial. Values are  $\pm$ SEM,  $n = 3$  for the enrichment formulations,  $n = 4$  for larvae (50 larvae per tank) and  $n = 6$  for enriched rotifers. Letters indicate significant differences between treatments.

have played a role in the increase in survival from the Fa and Fb larvae. The Fc formulation appears to be adequate in providing the necessary amount and proportions of fatty acid to snook larvae in reference to the wild eggs; however, even though growth and survival were not different than that of Algamac 3050, the rate of functional swim bladder development was significantly lower. Additional studies should therefore investigate the use of ARA powder rather than oil as mentioned previously, or improve the stability of the oil emulsion. Bessonart *et al.* (1999) did not observe a significant improvement in growth after supplementing ARA to gilthead sea bream larvae for 2 weeks, but a difference was observed after 3 weeks. In addition, Koven *et al.* (2001) did not

observe growth or survival improvement either after supplementing larvae from the same species for 2 weeks; nevertheless, a significant increase in stress resistance was observed in the supplemented larvae exposed to handling stress potentially suggesting more robust larvae. Therefore, future studies should investigate the effect of an ARA supplementation over a longer period of time and include the study of additional parameters such as cortisol production.

In conclusion, these studies bring fundamental information on the early fatty acid requirements of pompano and snook larvae. Findings also provided the first insight of snook larvae fatty acid incorporation during the live food period and the influ-

ence of an ARA supplementation on final fatty acid incorporation and ratios in snook larvae.

## Acknowledgments

The authors thank the research team at the Mote Marine Laboratory Aquaculture Park, as well as Tim Reed from Reed Mariculture for donating the specifically formulated enrichments, and Cynthia Faulk from the Fisheries and Mariculture Laboratory at the University of Texas Marine Science Institute for her help with the lipid analysis. Funding for this study was provided by the University of Stirling, the Florida Department of Agriculture and Consumer Services, Adam H. Putnam, Commissioner, Contract No. 017138, Florida Fish and Wildlife Conservation Commission (Contract No.10087, Permit # SAL 09-522-SR) and the National Sea Grant College Program of the U.S. Department of Commerce's National Oceanic and Atmospheric Administration (NOAA), Grant No. NA100AR4170079. The views are those of the authors and do not necessarily reflect the view of these organizations.

## References

- Atalah E., Hernández-Cruz C.M., Ganuza E., Benítez-Santana T., Ganga R., Roo J., Montero D. & Izquierdo M. (2011) Importance of dietary arachidonic acid for the growth, survival and stress resistance of larval European sea bass (*Dicentrarchus labrax*) fed high dietary docosahexaenoic and eicosapentaenoic acids. *Aquaculture Research* **42**, 1261–1268. doi: 10.1111/j.1365-2109.2010.02714.x.
- Atalah E., Maria Hernandez-Cruz C., Benitez-Santana T., Ganga R., Roo J. & Izquierdo M. (2011) Importance of the relative levels of dietary arachidonic acid and eicosapentaenoic acid for culture performance of gilthead seabream (*Sparus aurata*) larvae. *Aquaculture Research* **42**, 1279–1288. doi: 10.1111/j.1365-2109.2010.02716.x.
- Bell G. (2003) Arachidonic acid in aquaculture feeds: current status and future opportunities. *Aquaculture* **218**, 491–499. doi: 10.1016/S0044-8486(02)00370-8.
- Bell J.G., Castell J.D., Tocher D.R., Macdonald F.M. & Sargent J.R. (1995) Effects of different dietary arachidonic acid : docosahexaenoic acid ratios on phospholipid fatty acid compositions and prostaglandin production in juvenile turbot (*Scophthalmus maximus*). *Fish Physiology and Biochemistry* **14**, 139–151. doi: 10.1007/BF00002457.
- Bell M.V., Batty R.S., Dick J.R., Fretwell K., Navarro J.C. & Sargent J.R. (1995) Dietary deficiency of docosahexaenoic acid impairs vision at low light intensities in juvenile herring (*Clupea harengus* L.). *Lipids* **30**, 443–449.
- Bell J.G., Ashton I., Secombes C.J., Weitzel B.R., Dick J.R. & Sargent J.R. (1996) Dietary lipid affects phospholipid fatty acid compositions, eicosanoid production and immune function in Atlantic salmon (*Salmo salar*). *Prostaglandins, Leukotrienes and Essential Fatty Acids* **54**, 173–182.
- Benítez-Santana T., Masuda R., Juárez Carrillo E., Ganuza E., Valencia A., Hernández-Cruz C.M. & Izquierdo M.S. (2007) Dietary n-3 HUFA deficiency induces a reduced visual response in gilthead seabream *Sparus aurata* larvae. *Aquaculture* **264**, 408–417. doi: 10.1016/j.aquaculture.2006.10.024
- Bessonart M., Izquierdo M., Salhi M., Hernández-Cruz C., González M. & Fernández-Palacios H. (1999) Effect of dietary arachidonic acid levels on growth and survival of gilthead sea bream (*Sparus aurata* L.) larvae. *Aquaculture* **179**, 265–275.
- Cahu C., Zambonino Infante J. & Takeuchi T. (2003) Nutritional components affecting skeletal development in fish larvae. *Aquaculture* **227**, 245–258. doi: 10.1016/S0044-8486(03)00507-6.
- Castell J.D., Bell J.G., Tocher D.R. & Sargent J.R. (1994) Effects of purified diets containing different combinations of arachidonic and docosahexaenoic acid on survival, growth and fatty acid composition of juvenile turbot (*Scophthalmus maximus*). *Aquaculture* **128**, 315–333.
- Cejas J.R., Almansa E., Jérez S., Bolaños A., Felipe B. & Lorenzo A. (2004) Changes in lipid class and fatty acid composition during development in white seabream (*Diplodus sargus*) eggs and larvae. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* **139**, 209–216. doi: 10.1016/j.cbpc.2004.07.010.
- Conceição L.E.C., Yúfera M., Makridis P., Morais S. & Dinis M.T. (2010) Live feeds for early stages of fish rearing. *Aquaculture Research* **41**, 613–640. doi: 10.1111/j.1365-2109.2009.02242.x.
- Cunha I., Galante-Oliveira S., Rocha E., Planas M., Urbatzka R. & Castro L.F.C. (2013) Dynamics of PPARs, fatty acid metabolism genes and lipid classes in eggs and early larvae of a teleost. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* **164**, 247–258.
- Estevez A., Kaneko T., Seikai T., Tagawa M. & Tanaka M. (2001) ACTH and MSH production in Japanese flounder (*Paralichthys olivaceus*) larvae fed arachidonic acid-enriched live prey. *Aquaculture* **192**, 309–319.
- Faulk C.K. & Holt G.J. (2003) Lipid nutrition and feeding of cobia *Rachycentron canadum* larvae. *Journal of the World Aquaculture Society* **34**, 368–378.
- Faulk C. & Holt G.J. (2005) Advances in rearing cobia larvae in recirculating aquaculture systems: live prey enrichment and greenwater culture. *Aquaculture* **249**, 231–243. doi: 10.1016/j.aquaculture.2005.03.033.
- Fogerty A.C., Evans A.J., Ford G.L. & Kennett B.H. (1986) Distribution of omega 6 and omega 3 fatty

- acids in lipid classes in Australian fish. *Nutrition Reports International* **33**, 777–786.
- Folch J., Lees M. & Sloane Stanley G.H. (1957) A simple method for the isolation and purification of total lipides from animal tissues. *Journal of Biological Chemistry* **226**, 497–509.
- Ganga R., Tort L., Acerete L., Montero D. & Izquierdo M.S. (2006) Modulation of ACTH-induced cortisol release by polyunsaturated fatty acids in interrenal cells from gilthead seabream, *Sparus aurata*. *Journal of Endocrinology* **190**, 39–45.
- Gibson R.A. (1983) Australian fish—An excellent source of both arachidonic acid and  $\omega$ -3 polyunsaturated fatty acids. *Lipids* **18**, 743–752. doi: 10.1007/BF02534631.
- Glencross B.D. (2009) Exploring the nutritional demand for essential fatty acids by aquaculture species. *Reviews in Aquaculture* **1**, 71–124. doi: 10.1111/j.1753-5131.2009.01006.x.
- Hamre K., Yúfera M., Rønnestad I., Boglione C., Conceição L.E.C. & Izquierdo M. (2013) Fish larval nutrition and feed formulation: knowledge gaps and bottlenecks for advances in larval rearing. *Reviews in Aquaculture* **5**, S26–S58. doi: 10.1111/j.1753-5131.2012.01086.x.
- Harel M., Gavasso S., Leshin J., Gubernatis A. & Place A.R. (2001) The effect of tissue docosahexaenoic and arachidonic acids levels on hypersaline tolerance and leucocyte composition in striped bass (*Morone saxatilis*) larvae. *Fish Physiology and Biochemistry* **24**, 113–123. doi: 10.1023/A:1011924704459.
- Heming T. & Buddington R. (1988) Yolk absorption in embryonic and larval fishes. In: *Fish Physiology* (ed. by W. Hoar & D. Randall), pp. 407–446. Elsevier, London.
- Hunter J.R. (1981) Feeding ecology and predation of marine fish larvae. In: *Marine Fish Larvae Morphology Ecology and Relation to Fisheries* (ed. by R. Lasker), pp. 33–77. University of Washington Press, Seattle, WA.
- Izquierdo M.S. (1996) Review Article Essential fatty acid requirements of cultured marine fish larvae. *Aquaculture Nutrition* **2**, 183–191.
- Izquierdo M.S., Socorro J., Arantzamendi L. & Hernández-Cruz C.M. (2000) Recent advances in lipid nutrition in fish larvae. *Fish Physiology and Biochemistry* **22**, 97–107. doi: 10.1023/a:1007810506259.
- Kamler E. (2007) Resource allocation in yolk-feeding fish. *Reviews in Fish Biology and Fisheries* **18**, 143–200. doi: 10.1007/s11160-007-9070-x.
- Koven W.M., Kissil G.W. & Tandler A. (1989) Lipid and n-3 requirement of *Sparus aurata* larvae during starvation and feeding. *Aquaculture* **79**, 185–191.
- Koven W., Barr Y., Lutzky S., Ben-Atia I., Weiss R., Harel M., Behrens P. & Tandler A. (2001) The effect of dietary arachidonic acid (20:4n-6) on growth, survival and resistance to handling stress in gilthead seabream (*Sparus aurata*) larvae. *Aquaculture* **193**, 107–122.
- Main K.L., Resley M.J., Rhody N.R., Nystrom M., Stevens T. & Adams C., 2010. An overview of Florida pompano (*Trachinotus carolinus*) research at Mote Aquaculture Research Park. *Aquaculture America Conference San Diego*. Available at <https://www.was.org/meetings/SessionAbstracts.aspx?code=aq2010&session=28>.
- McEvoy L.A., Estevez A., Bell J.G., Shields R.J., Gara B. & Sargent J.R. (1998) Influence of dietary levels of eicosapentaenoic and arachidonic acids on the pigmentation success of turbot (*Scophthalmus maximus*) and halibut (*Hippoglossus hippoglossus*). *Bulletin-Aquaculture Association of Canada* **98**, 17–20.
- Montero D., Kalinowski T., Obach A., Robaina L., Tort L., Caballero M. & Izquierdo M. (2003) Vegetable lipid sources for gilthead seabream (*Sparus aurata*): effects on fish health. *Aquaculture* **225**, 353–370. doi: 10.1016/S0044-8486(03)00301-6
- Morais S., Mourente G., Ortega A., Tocher J.A. & Tocher D.R. (2011) Expression of fatty acyl desaturase and elongase genes, and evolution of DHA:EPA ratio during development of unfed larvae of Atlantic bluefin tuna (*Thunnus thynnus* L.). *Aquaculture* **313**, 129–139.
- Morrison W.R. & Smith L.M. (1964) Preparation of fatty acid methyl esters and dimethylacetals from lipids with boron fluoride-methanol. *Journal of Lipid Research* **5**, 600–608.
- Mourente G. & Vázquez R. (1996) Changes in the content of total lipid, lipid classes and their fatty acids of developing eggs and unfed larvae of the Senegal sole, *Solea senegalensis* Kaup. *Fish Physiology and Biochemistry* **15**, 221–235. doi: 10.1007/BF01875573.
- Mourente G., Rodriguez A., Tocher D.R. & Sargent J.R. (1993) Effects of dietary docosahexaenoic acid (DHA; 22: 6n-3) on lipid and fatty acid compositions and growth in gilthead sea bream (*Sparus aurata* L.) larvae during first feeding. *Aquaculture* **112**, 79–98.
- Mourente G., Rodriguez A., Grau A. & Pastor E. (1999) Utilization of lipids by *Dentex dentex* L. (Osteichthyes, Sparidae) larvae during lecithotrophia and subsequent starvation. *Fish Physiology and Biochemistry* **21**, 45–58.
- Neuringer M., Anderson G.J. & Connor W.E. (1988) The essentiality of n-3 fatty acids for the development and function of the retina and brain. *Annual Review of Nutrition* **8**, 517–541. doi: 10.1146/annurev.nu.08.070188.002505.
- Ogata H.Y., Emata A.C., Garibay E.S. & Furuita H. (2004) Fatty acid composition of five candidate aquaculture species in Central Philippines. *Aquaculture* **236**, 361–375. doi: 10.1016/j.aquaculture.2003.10.015.
- Rainuzzo J.R., Reitan K.I. & Jørgensen L. (1992) Comparative study on the fatty-acid and lipid composition of 4 marine fish larvae. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* **103**, 21–26. doi: 10.1016/0305-0491(92)90407-I.
- Rainuzzo J.R., Reitan K.I., Jørgensen L., Olsen Y. & Jørgensen L. (1994) Lipid composition in turbot larvae fed live feed cultured by emulsions of different lipid classes. *Comparative Biochemistry and Physiology Part A: Physiology* **107**, 699–710.

- Rodriguez C., Pérez J.A., Badía P., Izquierdo M.S., Fernández-Palacios H. & Lorenzo Hernandez A. (1998) The n-3 highly unsaturated fatty acids requirements of gilthead seabream (*Sparus aurata* L.) larvae when using an appropriate DHA EPA ratio in the diet. *Aquaculture* **169**, 9–23.
- Rønnestad I. (1999) Fish larval nutrition: a review of recent advances in the roles of amino acids. *Aquaculture* **177**, 201–216. doi: 10.1016/S0044-8486(99)00082-4.
- Rønnestad I., Finn R.N., Lein I. & Lie O. (1995) Compartmental changes in the contents of total lipid, lipid classes and their associated fatty acids in developing yolk-sac larvae of Atlantic halibut, *Hippoglossus hippoglossus* (L.). *Aquaculture Nutrition* **1**, 119–130.
- Rønnestad I., Koven W.M., Tandler A., Harel M. & Fyhn H.J. (1994) Energy metabolism during development of eggs and larvae of gilthead sea bream (*Sparus aurata*). *Marine Biology* **120**, 187–196. doi: 10.1007/BF00349678.
- Roo F.J., Hernández-Cruz C.M., Socorro J.A., Fernández-Palacios H., Montero D. & Izquierdo M.S. (2009) Effect of DHA content in rotifers on the occurrence of skeletal deformities in red porgy *Pagrus pagrus* (Linnaeus, 1758). *Aquaculture* **287**, 84–93. doi: 10.1016/j.aquaculture.2008.10.010.
- Sargent J.R., Henderson R.J. & Tocher D.R. (1989) The lipids. In: *Fish Nutrition* (ed. by J.E. Halver), pp. 153–218. Interscience Publishers, New York.
- Sargent J.R., McEvoy L.A. & Bell G. (1997) Requirements, presentation and sources of polyunsaturated fatty acids in marine fish larval feeds. *Aquaculture* **155**, 117–127. doi: 10.1016/S0044-8486(97)00122-1.
- Sargent J.R., Bell G., McEvoy L.A., Tocher D.R. & Estévez A. (1999) Recent developments in the essential fatty acid nutrition of fish. *Aquaculture* **177**, 191–199. doi: 10.1016/S0044-8486(99)00083-6.
- Sargent J.R., McEvoy L., Estévez A., Bell G., Bell M.V., Henderson R.J. & Tocher D.R. (1999) Lipid nutrition of marine fish during early development: current status and future directions. *Aquaculture* **179**, 217–229. doi: 10.1016/S0044-8486(99)00191-X.
- Sargent J.R., Tocher D.R. & Bell G. (2002) The lipids. In: *Fish Nutrition* (ed. by J. Halver & R. Hardy), pp. 181–257. Elsevier, San Diego.
- Stottrup J.G. (2000) The elusive copepods: their production and suitability in marine aquaculture. *Aquaculture Research* **31**, 703–711.
- Tandler A., Watanabe T., Satoh S. & Fukusho K. (1989) The effect of food deprivation on the fatty acid and lipid profile of red seabream (*Pagrus major*) larvae. *British Journal of Nutrition* **62**, 349–361.
- Tocher D.R. (1995) Glycerophospholipid metabolism. In: *Biochemistry and Molecular Biology of Fishes* (ed. by P.W. Hochachka & T.P. Mommsen), pp. 119–157. Elsevier Science B.V., Vancouver, BC.
- Tocher D.R. (2003) Metabolism and functions of lipids and fatty acids in teleost fish. *Reviews in Fisheries Science* **11**, 107–184.
- Tocher D.R., Fraser A.J., Sargent J.R. & Gamble J.C. (1985a) Lipid class composition during embryonic and early larval development in Atlantic herring (*Clupea harengus*, L.). *Lipids* **20**, 84–89. doi: 10.1007/BF02534213.
- Tocher D.R., Fraser A.J., Sargent J.R. & Gamble J.C. (1985b) Fatty acid composition of phospholipids and neutral lipids during embryonic and early larval development in Atlantic herring (*Clupea harengus*, L.). *Lipids* **20**, 69–74. doi: 10.1007/BF02534210.
- Tocher D.R., Bell J.G. & Sargent J.R. (1996) Production of eicosanoids derived from 20:4n-6 and 20:5n-3 in primary cultures of turbot (*Scophthalmus maximus*) brain astrocytes in response to platelet activating factor, substance P and interleukin-1 beta. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* **115**, 215–222.
- Van der Meeren T., Wilhelmssen S., Klungsosyr J. & Kvenseth P.G. (1993) Fatty acid composition of unfed cod larvae *Gadus morhua* L. and cod larvae feeding on natural plankton in large enclosures. *Journal of the World Aquaculture Society* **24**, 167–185. doi: 10.1111/j.1749-7345.1993.tb00006.x.
- Van der Meeren T., Olsen R.E., Hamre K. & Fyhn H.J. (2008) Biochemical composition of copepods for evaluation of feed quality in production of juvenile marine fish. *Aquaculture* **274**, 375–397. doi: 10.1016/j.aquaculture.2007.11.041.
- Vetter R.D., Hodson R.E. & Arnold C. (1983) Energy metabolism in a rapidly developing marine fish egg, the red drum (*Sciaenops ocellata*). *Canadian Journal of Fisheries and Aquatic Sciences* **40**, 627–634.
- Villalta M., Estévez A., Bransden M.P. & Bell J.G. (2008) Arachidonic acid, arachidonic/eicosapentaenoic acid ratio, stearidonic acid and eicosanoids are involved in dietary-induced albinism in Senegal sole (*Solea senegalensis*). *Aquaculture Nutrition* **14**, 120–128. doi: 10.1111/j.1365-2095.2007.00511.x.
- Watanabe T. (1993) Importance of docosahexaenoic acid in marine larval fish. *Journal of the World Aquaculture Society* **24**, 152–161. doi: 10.1111/j.1749-7345.1993.tb00004.x.
- Watanabe T., Kitajima C. & Fujita S. (1983) Nutritional values of live organisms used in Japan for mass propagation of fish: a review. *Aquaculture* **34**, 115–143.
- Yanes-Roca C., Rhody N., Nystrom M. & Main K.L. (2009) Effects of fatty acid composition and spawning season patterns on egg quality and larval survival in common snook (*Centropomus undecimalis*). *Aquaculture* **287**, 335–340. doi: 10.1016/j.aquaculture.2008.10.043.